

Direct phase-sensitive identification of a d -form factor density wave in underdoped cuprates

Kazuhiro Fujita^{a,b,c,1}, Mohammad H. Hamidian^{a,b,1}, Stephen D. Edkins^{b,d}, Chung Koo Kim^a, Yuhki Kohsaka^e, Masaki Azuma^f, Mikio Takano^g, Hidenori Takagi^{c,h,i}, Hiroshi Eisaki^j, Shin-ichi Uchida^c, Andrea Allais^k, Michael J. Lawler^{b,l}, Eun-Ah Kim^b, Subir Sachdev^{k,m}, and J. C. Séamus Davis^{a,b,d,2}

^aCondensed Matter Physics and Materials Science Department, Brookhaven National Laboratory, Upton, NY 11973; ^bLaboratory of Atomic and Solid State Physics, Department of Physics, Cornell University, Ithaca, NY 14853; ^cDepartment of Physics, University of Tokyo, Bunkyo-ku, Tokyo 113-0033, Japan; ^dSchool of Physics and Astronomy, University of St. Andrews, Fife KY16 9SS, Scotland; ^eRIKEN Center for Emergent Matter Science, Wako, Saitama 351-0198, Japan; ^fMaterials and Structures Laboratory, Tokyo Institute of Technology, Yokohama, Kanagawa 226-8503, Japan; ^gInstitute for Integrated Cell-Material Sciences, Kyoto University, Sakyo-ku, Kyoto 606-8501, Japan; ^hRIKEN Advanced Science Institute, Wako, Saitama 351-0198, Japan; ⁱMax-Planck-Institut für Festkörperforschung, 70569 Stuttgart, Germany; ^jNanoelectronics Research Institute, National Institute of Advanced Industrial Science and Technology, Tsukuba, Ibaraki 305-8568, Japan; ^kDepartment of Physics, Harvard University, Cambridge, MA 02138; ^lDepartment of Physics and Astronomy, Binghamton University, Binghamton, NY 13902; and ^mPerimeter Institute for Theoretical Physics, Waterloo, ON, Canada N2L 2Y5

Contributed by J. C. Séamus Davis, May 30, 2014 (sent for review April 1, 2014; reviewed by Joe Orenstein and Vidya Madhavan)

The identity of the fundamental broken symmetry (if any) in the underdoped cuprates is unresolved. However, evidence has been accumulating that this state may be an unconventional density wave. Here we carry out site-specific measurements within each CuO₂ unit cell, segregating the results into three separate electronic structure images containing only the Cu sites [Cu(*r*)] and only the *x*/*y* axis O sites [O_{*x*}(*r*) and O_{*y*}(*r*)]. Phase-resolved Fourier analysis reveals directly that the modulations in the O_{*x*}(*r*) and O_{*y*}(*r*) sublattice images consistently exhibit a relative phase of π . We confirm this discovery on two highly distinct cuprate compounds, ruling out tunnel matrix-element and materials-specific systematics. These observations demonstrate by direct sublattice phase-resolved visualization that the density wave found in underdoped cuprates consists of modulations of the intraunit-cell states that exhibit a predominantly d -symmetry form factor.

broken symmetries weaken simultaneously with increasing p and disappear jointly near $p_c = 0.19$ (13). For multiple cuprate compounds, neutron scattering reveals clear intraunit-cell breaking of rotational symmetry (14, 15, 16). Thermal transport studies (17) can likewise be interpreted. Polarized X-ray scattering studies reveal the electronic inequivalence between O_{*x*} and O_{*y*} sites (18) and that angular dependent scattering is best modeled by spatially modulating their inequivalence with a d -symmetry form factor (19). Thus, evidence from a variety of techniques indicates that $Q = 0$ C₄ breaking (electronic inequivalence of O_{*x*} and O_{*y*}) is a key element of underdoped-cuprate electronic structure. The apparently distinct phenomenology of $Q \neq 0$ incommensurate density waves (DW) in underdoped cuprates has also been reported extensively (20–27). Moreover, recent studies (28, 29) have demonstrated beautifully that the density modulations first visualized by scanning tunneling microscopy (STM) imaging (20) are in-