

# Holography of compressible quantum phases

KITP, September 29, 2011

[sachdev.physics.harvard.edu](mailto:sachdev.physics.harvard.edu)





**Rob Myers**



**Ajay Singh**



**Liza Huijse**



**Matthias Punk**



**Max Metlitski**



**Sean Hartnoll**


Conformal quantum matter

Compressible quantum matter

Conformal quantum matter


Compressible quantum matter

# Superfluid-insulator transition



Ultracold  $^{87}\text{Rb}$   
atoms - bosons

M. Greiner, O. Mandel, T. Esslinger, T. W. Hänsch, and I. Bloch, *Nature* **415**, 39 (2002).



# Quantum critical transport

Quantum “*nearly perfect fluid*”  
with shortest possible  
equilibration time,  $\tau_{\text{eq}}$

$$\tau_{\text{eq}} = \mathcal{C} \frac{\hbar}{k_B T}$$

where  $\mathcal{C}$  is a *universal* constant

# AdS<sub>4</sub> theory of “nearly perfect fluids”

To leading order in a gradient expansion, charge transport in an infinite set of strongly-interacting CFT3s can be described by Einstein-Maxwell gravity/electrodynamics on AdS<sub>4</sub>-Schwarzschild


We include all possible 4-derivative terms: after suitable field redefinitions, the required theory has only *one* dimensionless constant  $\gamma$  ( $L$  is the radius of AdS<sub>4</sub>):

$$\mathcal{S}_{EM} = \int d^4x \sqrt{-g} \left[ -\frac{1}{4e^2} F_{ab} F^{ab} + \frac{\gamma L^2}{e^2} C_{abcd} F^{ab} F^{cd} \right],$$

where  $C_{abcd}$  is the Weyl curvature tensor.


*Stability and causality constraints restrict  $|\gamma| < 1/12$ .*

# AdS<sub>4</sub> theory of strongly interacting “perfect fluids”




R. C. Myers, S. Sachdev, and A. Singh, *Physical Review D* **83**, 066017 (2011)

# AdS<sub>4</sub> theory of strongly interacting “perfect fluids”




R. C. Myers, S. Sachdev, and A. Singh, *Physical Review D* **83**, 066017 (2011)

# AdS<sub>4</sub> theory of strongly interacting “perfect fluids”




R. C. Myers, S. Sachdev, and A. Singh, *Physical Review D* **83**, 066017 (2011)

# AdS<sub>4</sub> theory of strongly interacting “perfect fluids”



R. C. Myers, S. Sachdev, and A. Singh, *Physical Review D* **83**, 066017 (2011)

# AdS<sub>4</sub> theory of strongly interacting “perfect fluids”



R. C. Myers, S. Sachdev, and A. Singh, *Physical Review D* **83**, 066017 (2011)

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories: the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on the heavy fermion compounds and the cuprate superconductors

# Compressible quantum matter

- Consider an infinite, continuum, translationally-invariant quantum system with a globally conserved U(1) charge  $Q$  (the “electron density”) in spatial dimension  $d > 1$ .

# Compressible quantum matter

- Consider an infinite, continuum, translationally-invariant quantum system with a globally conserved U(1) charge  $Q$  (the “electron density”) in spatial dimension  $d > 1$ .
- Describe zero temperature phases where  $d\langle Q \rangle / d\mu \neq 0$ , where  $\mu$  (the “chemical potential”) which changes the Hamiltonian,  $H$ , to  $H - \mu Q$ .

# Compressible quantum matter

- Consider an infinite, continuum, translationally-invariant quantum system with a globally conserved U(1) charge  $Q$  (the “electron density”) in spatial dimension  $d > 1$ .
- Describe zero temperature phases where  $d\langle Q \rangle / d\mu \neq 0$ , where  $\mu$  (the “chemical potential”) which changes the Hamiltonian,  $H$ , to  $H - \mu Q$ .
- Compressible systems must be gapless.


# Compressible quantum matter

- Consider an infinite, continuum, translationally-invariant quantum system with a globally conserved U(1) charge  $Q$  (the “electron density”) in spatial dimension  $d > 1$ .
- Describe zero temperature phases where  $d\langle Q \rangle/d\mu \neq 0$ , where  $\mu$  (the “chemical potential”) which changes the Hamiltonian,  $H$ , to  $H - \mu Q$ .
- Compressible systems must be gapless.
- Conformal systems are compressible in  $d = 1$ , but not for  $d > 1$ .

# Compressible quantum matter

One compressible state is the **solid** (or “Wigner crystal” or “stripe”).

This state breaks translational symmetry.



# Compressible quantum matter


Another familiar compressible state is  
the **superfluid**.

This state breaks the global  $U(1)$   
symmetry associated with  $Q$



Condensate of  
fermion pairs

The only other familiar compressible phase is a **Fermi Liquid** with a **Fermi surface**



Graphene


# The Landau Fermi liquid

- The *only* low energy excitations are long-lived quasiparticles near the Fermi surface.
- **Luttinger relation:** The total “volume (area)”  $\mathcal{A}$  enclosed by the Fermi surface is equal to  $\langle Q \rangle$ . This is a *key* constraint which allows extrapolation from weak to strong coupling, and also holds for “non-Fermi liquid” compressible phases to be discussed later.




Area  $\mathcal{A} = \langle Q \rangle$

# Begin with a CFT




Dirac fermions + gauge field + .....

# Holographic representation: AdS<sub>4</sub>




$$\mathcal{S} = \int d^4x \sqrt{-g} \left[ \frac{1}{2\kappa^2} \left( R + \frac{6}{L^2} \right) \right]$$

Will describe a Landau Fermi liquid  
obtained by applying a chemical potential to  
the “deconfined” CFT




# The Maxwell-Einstein theory of the applied chemical potential yields a AdS<sub>4</sub>-Reissner-Nordström black-brane



$$\mathcal{S} = \int d^4x \sqrt{-g} \left[ \frac{1}{2\kappa^2} \left( R + \frac{6}{L^2} \right) - \frac{1}{4e^2} F_{ab} F^{ab} \right]$$


# Holographic theory of a Fermi liquid

S. Sachdev  
arXiv:1107.5321



$$\mathcal{S} = \int d^4x \sqrt{-g} \left[ \frac{1}{2\kappa^2} \left( R + \frac{6}{L^2} \right) - \frac{Z(\phi)}{4e^2} F_{ab} F^{ab} + \mathcal{L}[\text{matter}, \phi] \right]$$

In a confining phase, the horizon disappears, there is charge density delocalized in the bulk spacetime, and a Fermi liquid phase is obtained on the boundary



Gauss Law in the bulk

$\Leftrightarrow$  Luttinger theorem on the boundary

In a confining phase, the horizon disappears,  
there is charge density delocalized in the bulk spacetime,  
and a Fermi liquid phase is obtained on the boundary

Consider QED<sub>4</sub>, with *full* quantum fluctuations,

$$S = \int d^4x \sqrt{g} \left[ \frac{1}{4e^2} F_{ab} F^{ab} + i (\bar{\psi} \Gamma^M D_M \psi + m \bar{\psi} \psi) \right].$$


in a metric which is AdS<sub>4</sub> in the UV, and confining in the IR.  
A simple model

$$ds^2 = \frac{1}{z^2} (dz^2 - dt^2 + dx^2 + dy^2) \quad , \quad z < z_m$$

with  $z_m$  determined by the confining scale.

# Holographic theory of a Fermi liquid

S. Sachdev  
arXiv:1107.5321




Massive Dirac fermions at zero chemical potential

$$\text{Dispersion } E_\ell(k) = \sqrt{k^2 + M_\ell^2}$$

$$\text{Masses } M_\ell \sim 1/z_m$$

# Holographic theory of a Fermi liquid

S. Sachdev  
arXiv:1107.5321




Massive Dirac fermions at zero chemical potential

$$\text{Dispersion } E_\ell(k) = \sqrt{k^2 + M_\ell^2}$$

$$\text{Masses } M_\ell \sim 1/z_m$$

Almost all previous holographic theories have considered the situation where the spacing between the  $E_\ell(k)$  vanishes, and an infinite number of  $E_\ell(k)$  are relevant.



The spectrum at non-zero chemical potential is determined by self-consistently solving the Dirac equation and Gauss's law:

$$\left(\vec{\Gamma} \cdot \vec{D} + m\right) \Psi_\ell = E_\ell \Psi_\ell ; \quad \nabla_z \mathcal{E}_z = \sum_\ell \int \frac{d^2 k}{4\pi^2} \Psi_\ell^\dagger(k, z) \Psi_\ell(k, z) f(E_\ell(k))$$

where  $\mathcal{E}$  is the electric field, and  $f(E)$  is the Fermi function

# Holographic theory of a Fermi liquid

S. Sachdev  
arXiv:1107.5321

- The confining geometry implies that all gauge and graviton modes are gapped (modulo Landau damping from the Fermi surface).

- The confining geometry implies that all gauge and graviton modes are gapped (modulo Landau damping from the Fermi surface).
- We can apply standard many body theory results, treating this multi-band system in 2 dimensions, like a 2DEG at a semiconductor surface.

- The confining geometry implies that all gauge and graviton modes are gapped (modulo Landau damping from the Fermi surface).
- We can apply standard many body theory results, treating this multi-band system in 2 dimensions, like a 2DEG at a semiconductor surface.
- Integrating Gauss's Law, we obtain

$$\mathcal{E}_z(\text{boundary}) - \mathcal{E}_z(\text{IR}) = \mathcal{A}$$

But  $\mathcal{E}_z(\text{boundary}) = \langle \mathcal{Q} \rangle$ , but the rules of AdS/CFT. So we obtain the usual Luttinger theorem of a Landau Fermi liquid,


$$\mathcal{A} = \langle \mathcal{Q} \rangle$$

*provided*  $\mathcal{E}_z(\text{IR}) = 0$ .

Technical notes:

- No source term is included at the boundary for the fermions
- The boundary fermion Green's function is computed by taking a suitable limit of the bulk Green's function (Klebanov, Witten):

$$G(r, r') = \lim_{z, z' \rightarrow 0} (zz')^\alpha G_B(r, z; r', z')$$



Gauss Law in the bulk

$\Leftrightarrow$  Luttinger theorem on the boundary

In a confining FL phase, the metric terminates, the bulk charge equals the boundary charge, and the electric flux vanishes in the IR.

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# The Fermi surface

This is the locus of zero energy singularities in momentum space in the two-point correlator of fermions carrying charge  $Q$ .

$$G_{\text{fermion}}^{-1}(k = k_F, \omega = 0) = 0.$$

Excitations with  $k < k_F$  are ‘hole’-like (negative energy), and those with  $k > k_F$  are ‘particle’-like (positive energy), or vice-versa.

# The Fermi surface

This is the locus of zero energy singularities in momentum space in the two-point correlator of fermions carrying charge  $Q$ .

$$G_{\text{fermion}}^{-1}(k = k_F, \omega = 0) = 0.$$

Excitations with  $k < k_F$  are ‘hole’-like (negative energy), and those with  $k > k_F$  are ‘particle’-like (positive energy), or vice-versa.

Boson Green’s functions can’t generically have such singularities because the negative energy bosons would Bose condense.

# The Fermi surface

This is the locus of zero energy singularities in momentum space in the two-point correlator of fermions carrying charge  $Q$ .

$$G_{\text{fermion}}^{-1}(k = k_F, \omega = 0) = 0.$$

Excitations with  $k < k_F$  are ‘hole’-like (negative energy), and those with  $k > k_F$  are ‘particle’-like (positive energy), or vice-versa.

Boson Green’s functions can’t generically have such singularities because the negative energy bosons would Bose condense.

**Luttinger relation:** Applies as long as the global U(1) symmetry associated with  $Q$  is unbroken. The total “volume (area)”  $\mathcal{A}$  enclosed by the Fermi surface is equal to  $\langle Q \rangle$ . Here  $\langle Q \rangle$  includes the charge carried by the bosons. This is a *key* constraint which allows extrapolation from weak to strong coupling.

Consider mixture of fermions  $f$  and bosons  $b$ .

$$\begin{aligned} \mathcal{L} &= f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f \\ &+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots \end{aligned}$$

Consider mixture of fermions  $f$  and bosons  $b$ .  
There is a  $U(1) \times U_b(1)$  symmetry  
and 2 conserved charges:

$$Q = f^\dagger f$$
$$Q_b = b^\dagger b$$

$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$
$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

Consider mixture of fermions  $f$  and bosons  $b$ .  
There is a  $U(1) \times U_b(1)$  symmetry  
and 2 conserved charges:

The 2 symmetries imply 2  
Luttinger constraints. How-  
ever, bosons at non-zero den-  
sity invariably Bose condense  
at  $T = 0$ , and so  $U_b(1)$  is  
broken. So there is only the  
single constraint on the  $f$  Fermi  
surface. This describes mix-  
tures of  $^3\text{He}$  and  $^4\text{He}$ .

$$Q = f^\dagger f$$
$$Q_b = b^\dagger b$$

$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$
$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

Consider mixture of fermions  $f$  and bosons  $b$ .  
There is a  $U(1) \times U_b(1)$  symmetry  
and 2 conserved charges:

$$\mathcal{A} = \langle \mathcal{Q} \rangle$$


$$\mathcal{Q} = f^\dagger f$$
$$\mathcal{Q}_b = b^\dagger b$$

Superfluid:  $\langle b \rangle \neq 0$

$U_b(1)$  broken;  $U(1)$  unbroken

$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$
$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

Increase the coupling  $g$  until the boson,  $b$ , and fermion,  $f$ , can bind into a ‘molecule’, the fermion  $c$ .



$$Q = f^\dagger f$$


$$Q_b = b^\dagger b$$

$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

Increase the coupling  $g$  until the boson,  $b$ , and fermion,  $f$ , can bind into a ‘molecule’, the fermion  $c$ .

Decouple the interaction between  $b$  and  $f$  by a fermion  $c$



$$Q = f^\dagger f$$



$$Q_b = b^\dagger b$$

$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 + \frac{1}{g} c^\dagger c - c^\dagger f b - c b^\dagger f^\dagger + \dots$$

In a phase with  $U_b(1)$  unbroken, there is a Luttinger relation for each conserved  $U(1)$  charge. However, the boson,  $b$  cannot have a Fermi surface in its Green's function, and so there is no area associated with it, although the boson density is *included* in the Luttinger relation

$$\begin{aligned} A_c + A_f &= \langle f^\dagger f \rangle = \langle Q \rangle \\ A_c &= \langle b^\dagger b \rangle = \langle Q_b \rangle \end{aligned}$$




The  $b$  bosons  
have bound  
with  $f$  fermions  
to form  $c$   
“molecules”



S. Powell, S. Sachdev, and H. P. Büchler, *Physical Review B* **72**, 024534 (2005)

P. Coleman, I. Paul, and J. Rech, *Physical Review B* **72**, 094430 (2005)

# Phase diagram of boson-fermion mixture



Superfluid:  $\langle b \rangle \neq 0$   
 $U_b(1)$  broken;  $U(1)$  unbroken



Normal:  $\langle b \rangle = 0$   
 $U(1) \times U_b(1)$  unbroken



$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

- Now gauge  $\mathcal{Q} - \mathcal{Q}_b$  by a dynamic gauge field  $A_a$ .  
This leaves fermion  $c$  gauge-invariant


$$\begin{aligned} \mathcal{L} &= f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f \\ &+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots \end{aligned}$$

- Now gauge  $\mathcal{Q} - \mathcal{Q}_b$  by a dynamic gauge field  $A_a$ .  
This leaves fermion  $c$  gauge-invariant



(Need a background neutralizing charge)

$$\begin{aligned} \mathcal{L} = & f^\dagger \left( \partial_\tau - iA_\tau - \frac{(\nabla - i\mathbf{A})^2}{2m} - \mu \right) f \\ & + b^\dagger \left( \partial_\tau + iA_\tau - \frac{(\nabla + i\mathbf{A})^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots \end{aligned}$$

# Phase diagram of boson-fermion mixture



Superfluid:  $\langle b \rangle \neq 0$   
 $U_b(1)$  broken;  $U(1)$  unbroken




Normal:  $\langle b \rangle = 0$   
 $U(1) \times U_b(1)$  unbroken





$$\mathcal{L} = f^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau - \frac{\nabla^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

# Phase diagram of U(1) gauge theory



Higgs/confining phase:  
Fermi liquid (FL)




Deconfined phase:  
Fractionalized  
Fermi liquid (FL\*)





$$\mathcal{L} = f^\dagger \left( \partial_\tau - iA_\tau - \frac{(\nabla - i\mathbf{A})^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau + iA_\tau - \frac{(\nabla + i\mathbf{A})^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

# Phase diagram of U(1) gauge theory



Higgs/confining phase:  
Fermi liquid (FL)



Deconfined phase:  
Fractionalized  
Fermi liquid (FL\*)



$$\mathcal{L} = f^\dagger \left( \partial_\tau - iA_\tau - \frac{(\nabla - i\mathbf{A})^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau + iA_\tau - \frac{(\nabla + i\mathbf{A})^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

## Phase diagram of U(1) gauge theory

- FL phase: Fermi surface of gauge-neutral fermions encloses total global charge  $Q$
- FL\* phase: Fermi surface of gauge neutral fermions encloses only part of the global charge  $Q$

Higgs/continuing phase:  
Fermi liquid (FL)


Fractionalized  
Fermi liquid (FL\*)

$s$

$$\mathcal{L} = f^\dagger \left( \partial_\tau - iA_\tau - \frac{(\nabla - i\mathbf{A})^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau + iA_\tau - \frac{(\nabla + i\mathbf{A})^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

# Phase diagram of U(1) gauge theory



Similar to theories obtained by adding a chemical potential to CFTs (with non-Abelian gauge fields) with known gravity duals

L. Huijse and S. Sachdev, arXiv:1104.5022



$$\mathcal{L} = f^\dagger \left( \partial_\tau - iA_\tau - \frac{(\nabla - i\mathbf{A})^2}{2m} - \mu \right) f$$

$$+ b^\dagger \left( \partial_\tau + iA_\tau - \frac{(\nabla + i\mathbf{A})^2}{2m_b} - \mu_b \right) b + s|b|^2 - g b^\dagger f^\dagger f b + \dots$$

# Compressible quantum matter

Conjecture: All compressible states which preserve translational and global  $U(1)$  symmetries must have FERM SURFACES, but they are not necessarily Fermi liquids.

- Such states obey the Luttinger relation

$$\sum_{\ell} q_{\ell} \mathcal{A}_{\ell} = \langle Q \rangle,$$

where the  $\ell$ 'th Fermi surface has fermionic quasiparticles with global  $U(1)$  charge  $q_{\ell}$  and encloses area  $\mathcal{A}_{\ell}$ .

- Non-Fermi liquids have quasiparticles coupled to deconfined gauge fields (or gapless bosonic modes at quantum critical points).

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases


1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors


# Holographic theory of a Fermi liquid

S. Sachdev  
arXiv:1107.5321



In a confining FL phase, the metric terminates, the bulk charge equals the boundary charge, and the electric flux vanishes in the IR.

# Holographic theory of a Fermi liquid



In a deconfined FL\* phase, the metric extends to infinity (representing critical IR modes), and part of the electric flux “leaks out”.

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)

## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach

3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# Compressible quantum matter

## Conventional phases

1. Holographic theory of the Fermi liquid (FL)


## Exotic phases

1. Continuum models with gauge theories:  
the fractionalized Fermi liquid (FL\*)

2. Holographic approach


3. Connections to models and experiments on  
the heavy fermion compounds and  
the cuprate superconductors

# Magnetic order and the heavy Fermi liquid in the Kondo lattice



$$\langle \vec{\varphi} \rangle \neq 0$$


Magnetic Metal:  
f-electron moments  
and  
c-conduction electron  
Fermi surface



$$\langle \vec{\varphi} \rangle = 0$$


Heavy Fermi liquid  
with “large” Fermi  
surface of  
hybridized f and  
c-conduction  
electrons

# Separating onset of SDW order and the heavy Fermi liquid in the Kondo lattice



$$\langle \vec{\varphi} \rangle \neq 0$$

Magnetic Metal:  
f-electron moments  
and  
c-conduction electron  
Fermi surface




$$\langle \vec{\varphi} \rangle = 0$$

Heavy Fermi liquid  
with “large” Fermi  
surface of  
hybridized f and  
c-conduction  
electrons




# Separating onset of SDW order and the heavy Fermi liquid in the Kondo lattice




$$\langle \vec{\varphi} \rangle \neq 0$$

Magnetic Metal:  
f-electron moments  
and  
c-conduction electron  
Fermi surface



$$\langle \vec{\varphi} \rangle = 0$$

Conduction electron  
Fermi surface  
and  
spin-liquid of  
f-electrons




$$\langle \vec{\varphi} \rangle = 0$$

Heavy Fermi liquid  
with "large" Fermi  
surface of  
hybridized f and  
c-conduction  
electrons




# Separating onset of SDW order and the heavy Fermi liquid in the Kondo lattice




$$\langle \vec{\varphi} \rangle \neq 0$$

Magnetic Metal:  
f-electron moments  
and  
c-conduction electron  
Fermi surface



$$\langle \vec{\varphi} \rangle = 0$$

Fractionalized Fermi  
liquid (FL\*) phase  
with no symmetry  
breaking and “small”  
Fermi surface




$$\langle \vec{\varphi} \rangle = 0$$

Heavy Fermi liquid  
with “large” Fermi  
surface of  
hybridized f and  
c-conduction  
electrons


T. Senthil, S. Sachdev, and M. Vojta, *Phys. Rev. Lett.* **90**, 216403 (2003)

# Separating onset of SDW order and the heavy Fermi liquid in the Kondo lattice



$$\langle \vec{\varphi} \rangle \neq 0$$


Magnetic Metal:  
 $f$ -electron moments  
 and  
 $c$ -conduction electron  
 Fermi surface



$$\langle \vec{\varphi} \rangle = 0$$

Fractionalized Fermi  
 liquid (FL\*) phase  
 with no symmetry  
 breaking and “small”  
 Fermi surface

Transition  
 described by  
 condensation of a  
 “slave boson” in a  
 spin liquid with  
 fermionic spinons




$$\langle \vec{\varphi} \rangle = 0$$

Heavy Fermi liquid  
 with “large” Fermi  
 surface of  
 hybridized  $f$  and  
 $c$ -conduction  
 electrons

T. Senthil, S. Sachdev, and M. Vojta, *Phys. Rev. Lett.* **90**, 216403 (2003)


# Separating onset of SDW order and the heavy Fermi liquid in the Kondo lattice




T. Senthil, S. Sachdev, and M. Vojta, *Phys. Rev. Lett.* **90**, 216403 (2003)

# Experimental perspective on same phase diagrams of Kondo lattice

J. Custers, P. Gegenwart,  
C. Geibel, F. Steglich,  
P. Coleman, and S. Paschen,  
*Phys. Rev. Lett.*  
**104**, 186402 (2010)







PHYSICAL REVIEW B **69**, 014415 (2004)

## Magnetic field induced non-Fermi-liquid behavior in YbAgGe single crystals

S. L. Bud'ko,<sup>1</sup> E. Morosan,<sup>1,2</sup> and P. C. Canfield<sup>1,2</sup>



J. Custers, P. Gegenwart, C. Geibel, F. Steglich, P. Coleman, and S. Paschen,  
*Phys. Rev. Lett.* **104**, 186402 (2010)




# Detaching the antiferromagnetic quantum critical point from the Fermi-surface reconstruction in YbRh<sub>2</sub>Si<sub>2</sub>

Nature Physics 5, 465 (2009)


S. Friedemann<sup>1\*</sup>, T. Westerkamp<sup>1</sup>, M. Brando<sup>1</sup>, N. Oeschler<sup>1</sup>, S. Wirth<sup>1</sup>, P. Gegenwart<sup>1,2</sup>, C. Krellner<sup>1</sup>, C. Geibel<sup>1</sup> and F. Steglich<sup>1\*</sup>

# Fermi surface reconstruction in a single band model



$$\langle \vec{\varphi} \rangle \neq 0$$

Metal with electron  
and hole pockets




$$\langle \vec{\varphi} \rangle = 0$$

Metal with “large”  
Fermi surface




# Separating onset of SDW order and Fermi surface reconstruction



$$\langle \vec{\varphi} \rangle \neq 0$$

Metal with electron  
and hole pockets




$$\langle \vec{\varphi} \rangle = 0$$

Metal with “large”  
Fermi surface



# Separating onset of SDW order and Fermi surface reconstruction




$$\langle \vec{\varphi} \rangle \neq 0$$

Metal with electron  
and hole pockets

Electron and/or hole  
Fermi pockets form in  
“local” SDW order, but  
quantum fluctuations  
destroy long-range  
SDW order

$$\langle \vec{\varphi} \rangle = 0$$

Fractionalized Fermi  
liquid (FL\*) phase  
with no symmetry  
breaking and “small”  
Fermi surface




$$\langle \vec{\varphi} \rangle = 0$$

Metal with “large”  
Fermi surface

Y. Qi and S. Sachdev, *Physical Review B* **81**, 115129 (2010); M. Punk and S. Sachdev, to appear;  
see also T. C. Ribeiro and X.-G. Wen, *Physical Review B* **74**, 155113 (2006)

# Separating onset of SDW order and Fermi surface reconstruction



$$\langle \vec{\varphi} \rangle \neq 0$$


Metal with electron  
and hole pockets

Transition  
described by  
condensation of  
bosonic spinons

Electron and/or hole  
Fermi pockets form in  
“local” SDW order, but  
quantum fluctuations  
destroy long-range  
SDW order

$$\langle \vec{\varphi} \rangle = 0$$


Fractionalized Fermi  
liquid (FL\*) phase  
with no symmetry  
breaking and “small”  
Fermi surface



$$\langle \vec{\varphi} \rangle = 0$$

Metal with “large”  
Fermi surface

Y. Qi and S. Sachdev, *Physical Review B* **81**, 115129 (2010); M. Punk and S. Sachdev, to appear;  
see also T. C. Ribeiro and X.-G. Wen, *Physical Review B* **74**, 155113 (2006)



## Reconstructed Fermi Surface of Underdoped $\text{Bi}_2\text{Sr}_2\text{CaCu}_2\text{O}_{8+\delta}$ Cuprate Superconductors

H.-B. Yang,<sup>1</sup> J. D. Rameau,<sup>1</sup> Z.-H. Pan,<sup>1</sup> G. D. Gu,<sup>1</sup> P. D. Johnson,<sup>1</sup> H. Claus,<sup>2</sup> D. G. Hinks,<sup>2</sup> and T. E. Kidd<sup>3</sup>

## Conclusions

# Quantum criticality and conformal field theories

- New insights and solvable models for diffusion and transport of strongly interacting systems near quantum critical points
- The description is far removed from, and complementary to, that of the quantum Boltzmann equation which builds on the quasiparticle/vortex picture.
- Prospects non-linear, and non-equilibrium transport

# Conclusions

## Compressible quantum matter

- Presented a holographic model of a Fermi liquid
- Fractionalized Fermi liquid (FL\*), appears in deconfined gauge theories, holographic models, and lattice theories of the heavy-fermion compounds and cuprates superconductors.
- Numerous plausible sightings of the FL\* phase in recent experiments