


Quantum Entanglement, Strange metals, and black holes




Subir Sachdev, Harvard University

Quantum entanglement

Quantum Entanglement: quantum superposition with more than one particle




Hydrogen molecule:




$$= \frac{1}{\sqrt{2}} (|\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle)$$


Quantum Entanglement: quantum superposition with more than one particle




Quantum Entanglement: quantum superposition with more than one particle



Quantum Entanglement: quantum superposition with more than one particle



Quantum Entanglement: quantum superposition with more than one particle



Quantum entanglement

**Quantum
entanglement**


**Black
holes**

Black Holes


Objects so dense that light is gravitationally bound to them.

In Einstein's theory, the region inside the black hole **horizon** is disconnected from the rest of the universe.

Horizon radius $R = \frac{2GM}{c^2}$





On September 14, 2015, LIGO detected the merger of two black holes, each weighing about 30 solar masses, with radii of about 100 km, 1.3 billion light years away



0.1 seconds later !






LIGO
September 14, 2015


Black Holes + Quantum theory

Around 1974, Bekenstein and Hawking showed that the application of the quantum theory across a black hole horizon led to many astonishing conclusions


Quantum Entanglement across a black hole horizon



Quantum Entanglement across a black hole horizon




Quantum Entanglement across a black hole horizon




Quantum Entanglement across a black hole horizon

There is long-range quantum entanglement between the inside and outside of a black hole



Quantum Entanglement across a black hole horizon

Hawking used this to show that black hole horizons have an entropy and a temperature
(because to an outside observer, the state of the electron inside the black hole is an unknown)




Quantum Entanglement across a black hole horizon



The Hawking temperature $k_B T_H = \frac{\hbar^2}{8\pi M \ell_P^2}$ and

the Bekenstein-Hawking (BH) black hole entropy $\frac{S}{k_B} = \frac{A}{4\ell_P^2}$

where $\ell_P = \sqrt{\hbar G/c^3}$ is the Planck length,
and A is the surface area of the black hole.

Note the entropy is proportional to the surface area
rather than the volume.





LIGO
September 14, 2015

- The Hawking temperature, T_H influences the radiation from the black hole at the very last stages of the ring-down (not observed so far). The ring-down (approach to thermal equilibrium) happens very rapidly in a time $\sim \frac{\hbar}{k_B T_H} = \frac{8\pi GM}{c^3} \sim 8$ milliseconds.

**Quantum
entanglement**


**Black
holes**

**Quantum
entanglement**

**Black
holes**

**Strange
metals**

High temperature superconductors





Power Efficiency/Capacity/Stability



Power Bottlenecks




Accommodate Renewable Power




Efficient Rotating Machines



Information Technology



Next Generation HEP



Ultra-High Magnetic Fields




Medical




Transport

High temperature superconductors



Antiferromagnet

Strange metal



Superconductor




Figure: K. Fujita and J. C. Seamus Davis

Antiferromagnet







Figure: K. Fujita and J. C. Seamus Davis

Square lattice of Cu sites




Square lattice of Cu sites



Remove density
 p electrons

Square lattice of Cu sites




Electrons entangle in (“Cooper”) pairs into chemical bonds

$$\text{[Diagram of a pair of teal circles in a blue oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

Superconductivity




Cooper pairs form quantum superpositions at different locations: “Bose-Einstein condensation” in which all pairs are “everywhere at the same time”

$$\text{Cooper pair} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

Superconductivity




Cooper pairs form quantum superpositions at different locations: “Bose-Einstein condensation” in which all pairs are “everywhere at the same time”

$$\text{Cooper pair} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

Superconductivity




Cooper pairs form quantum superpositions at different locations: “Bose-Einstein condensation” in which all pairs are “everywhere at the same time”

$$\text{Cooper pair} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

Superconductivity




Cooper pairs form quantum superpositions at different locations: “Bose-Einstein condensation” in which all pairs are “everywhere at the same time”

$$\text{Cooper pair} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{[Diagram of two sites in an oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{[Diagram of two sites in an oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{[Diagram of two sites in an oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{[Diagram of two sites in an oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{Diagram of two sites in an oval} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !




Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{Diagram of two sites in an oval} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Square lattice of Cu sites

High temperature superconductivity !



Electrons entangle by exchanging partners, and there is long-range quantum entanglement in the strange metal.

$$\text{[Diagram of two teal circles in a blue oval]} = |\uparrow\downarrow\rangle - |\downarrow\uparrow\rangle$$

Strange metal

Entangled electrons lead to “strange” temperature dependence of resistivity and other properties







Figure: K. Fujita and J. C. Seamus Davis

Almost all many-electron systems are described by the quasiparticle concept: a quasiparticle is an “excited lump” in the many-electron state which responds just like an ordinary particle.



real horse



quasi horse

Almost all many-electron systems are described by the quasiparticle concept: a quasiparticle is an “excited lump” in the many-electron state which responds just like an ordinary particle.

- **Quasiparticles are additive excitations:**

The low-lying excitations of the many-body system can be identified as a set $\{n_\alpha\}$ of quasiparticles with energy ε_α

$$E = \sum_{\alpha} n_{\alpha} \varepsilon_{\alpha} + \sum_{\alpha, \beta} F_{\alpha\beta} n_{\alpha} n_{\beta} + \dots$$

Almost all many-electron systems are described by the quasiparticle concept: a quasiparticle is an “excited lump” in the many-electron state which responds just like an ordinary particle.

- Quasiparticles eventually collide with each other. Such collisions eventually leads to thermal equilibration in a chaotic quantum state, but the equilibration takes a long time. In a Fermi liquid, this time is of order $\hbar E_F / (k_B T)^2$ as $T \rightarrow 0$, where E_F is the Fermi energy.

Quantum matter without quasiparticles

The complex quantum entanglement in the strange metal does not allow for any quasiparticle excitations.

Quantum matter without quasiparticles

The complex quantum entanglement in the strange metal does not allow for any quasiparticle excitations.

- Systems *without* quasiparticles, like the strange metal, reach quantum chaos much more quickly than those with quasiparticles.

Quantum matter without quasiparticles

The complex quantum entanglement in the strange metal does not allow for any quasiparticle excitations.

- Systems *without* quasiparticles, like the strange metal, reach quantum chaos much more quickly than those with quasiparticles.
- There is an *lower bound* on the phase coherence time (τ_φ), and the time to many-body quantum chaos (τ_L) in all many-body quantum systems as $T \rightarrow 0$:

$$\tau_\varphi \geq C \frac{\hbar}{k_B T} \quad (\text{SS, 1999})$$

$$\tau_L \geq \frac{\hbar}{2\pi k_B T} \quad (\text{Maldacena, Shenker, Stanford, 2015})$$

So *e.g.* we cannot have $\tau_\varphi \sim \hbar/\sqrt{Jk_B T}$ where J is a microscopic coupling.

Quantum matter without quasiparticles

The complex quantum entanglement in the strange metal does not allow for any quasiparticle excitations.

- Systems *without* quasiparticles, like the strange metal, reach quantum chaos much more quickly than those with quasiparticles.
- There is an *lower bound* on the phase coherence time (τ_φ), and the time to many-body quantum chaos (τ_L) in all many-body quantum systems as $T \rightarrow 0$:

$$\tau_\varphi \geq C \frac{\hbar}{k_B T} \quad (\text{SS, 1999})$$

$$\tau_L \geq \frac{\hbar}{2\pi k_B T} \quad (\text{Maldacena, Shenker, Stanford, 2015})$$

So *e.g.* we cannot have $\tau_\varphi \sim \hbar/\sqrt{Jk_B T}$ where J is a microscopic coupling.

- In the strange metal the inequalities become equalities as $T \rightarrow 0$, and the time $\hbar/(k_B T)$ influences numerous observables.

**Quantum
entanglement**

**Black
holes**

**Strange
metals**


**Quantum
entanglement**

**Black
holes**

**Strange
metals**


**A "toy model" which is both a
strange metal and a black hole!**

A simple model of a metal with quasiparticles




Pick a set of random positions

A simple model of a metal with quasiparticles




Place electrons randomly on some sites

A simple model of a metal with quasiparticles




Electrons move one-by-one randomly

A simple model of a metal with quasiparticles




Electrons move one-by-one randomly

A simple model of a metal with quasiparticles



Electrons move one-by-one randomly

A simple model of a metal with quasiparticles



Electrons move one-by-one randomly

A simple model of a metal with quasiparticles

$$H = \frac{1}{(N)^{1/2}} \sum_{i,j=1}^N t_{ij} c_i^\dagger c_j + \dots$$

$$c_i c_j + c_j c_i = 0 \quad , \quad c_i c_j^\dagger + c_j^\dagger c_i = \delta_{ij}$$


$$\frac{1}{N} \sum_i c_i^\dagger c_i = Q$$

t_{ij} are independent random variables with $\overline{t_{ij}} = 0$ and $\overline{|t_{ij}|^2} = t^2$


**Fermions occupying the eigenstates of a
 $N \times N$ random matrix**

A simple model of a metal with quasiparticles

Let ε_α be the eigenvalues of the matrix t_{ij}/\sqrt{N} . The fermions will occupy the lowest NQ eigenvalues, upto the Fermi energy E_F . The density of states is $\rho(\omega) = (1/N) \sum_\alpha \delta(\omega - \varepsilon_\alpha)$.



A simple model of a metal with quasiparticles



Many-body
level spacing
 $\sim 2^{-N}$


Quasiparticle
excitations with
spacing $\sim 1/N$

There are 2^N many
body levels with energy

$$E = \sum_{\alpha=1}^N n_{\alpha} \varepsilon_{\alpha},$$


where $n_{\alpha} = 0, 1$. Shown
are all values of E for a
single cluster of size
 $N = 12$. The ε_{α} have a
level spacing $\sim 1/N$.

The Sachdev-Ye-Kitaev (SYK) model




Pick a set of random positions

The Sachdev-Ye-Kitaev (SYK) model




Place electrons randomly on some sites

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model




Entangle electrons pairwise randomly

The Sachdev-Ye-Kitaev (SYK) model



The SYK model has “nothing but entanglement”

The Sachdev-Ye-Kitaev (SYK) model



This describes both a strange metal and a black hole!


The Sachdev-Ye-Kitaev (SYK) model

$$H = \frac{1}{(2N)^{3/2}} \sum_{i,j,k,\ell=1}^N J_{ij;kl} c_i^\dagger c_j^\dagger c_k c_\ell - \mu \sum_i c_i^\dagger c_i$$

$$c_i c_j + c_j c_i = 0 \quad , \quad c_i c_j^\dagger + c_j^\dagger c_i = \delta_{ij}$$

$$Q = \frac{1}{N} \sum_i c_i^\dagger c_i$$

$J_{ij;kl}$ are independent random variables with $\overline{J_{ij;kl}} = 0$ and $\overline{|J_{ij;kl}|^2} = J^2$
 $N \rightarrow \infty$ yields critical strange metal.



S. Sachdev and J. Ye, PRL **70**, 3339 (1993)

A. Kitaev, unpublished; S. Sachdev, PRX **5**, 041025 (2015)

The Sachdev-Ye-Kitaev (SYK) model

There are 2^N many body levels with energy E , which do not admit a quasiparticle decomposition. Shown are all values of E for a single cluster of size $N = 12$. The $T \rightarrow 0$ state has an entropy NS_0 with

$$S_0 = \frac{G}{\pi} + \frac{\ln(2)}{4} = 0.464848 \dots$$
$$< \ln 2$$


where G is Catalan's constant, for the half-filled case $Q = 1/2$.

GPS: A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)

Many-body level spacing $\sim 2^{-N} = e^{-N \ln 2}$

Non-quasiparticle excitations with spacing $\sim e^{-NS_0}$

A simple model of a metal with quasiparticles



Many-body
level spacing
 $\sim 2^{-N}$

Quasiparticle
excitations with
spacing $\sim 1/N$

There are 2^N many
body levels with energy

$$E = \sum_{\alpha=1}^N n_{\alpha} \varepsilon_{\alpha},$$

where $n_{\alpha} = 0, 1$. Shown
are all values of E for a
single cluster of size
 $N = 12$. The ε_{α} have a
level spacing $\sim 1/N$.

The Sachdev-Ye-Kitaev (SYK) model

There are 2^N many body levels with energy E , which do not admit a quasiparticle decomposition. Shown are all values of E for a single cluster of size $N = 12$. The $T \rightarrow 0$ state has an entropy NS_0 with

$$S_0 = \frac{G}{\pi} + \frac{\ln(2)}{4} = 0.464848 \dots$$
$$< \ln 2$$

where G is Catalan's constant, for the half-filled case $Q = 1/2$.

GPS: A. Georges, O. Parcollet, and S. Sachdev, PRB **63**, 134406 (2001)

Many-body level spacing $\sim 2^{-N} = e^{-N \ln 2}$

Non-quasiparticle excitations with spacing $\sim e^{-NS_0}$

SYK and black holes

- The SYK model has a non-zero GPS entropy, NS_0 as $T \rightarrow 0$.

A. Georges, O. Parcollet, and S. Sachdev,
PRB **63**, 134406 (2001)


- The SYK model has a phase-coherence time $\tau_\varphi \sim \hbar/(k_B T)$

O. Parcollet and A. Georges,
PRB **59**, 5341 (1999)

These properties indicate that SYK model ‘holographically’ realizes a black hole, and S_0 is the BH entropy.


S. Sachdev, PRL **105**, 151602 (2010)

SYK and black holes




SYK and black holes

S. Sachdev, PRL **105**, 151602 (2010)



The SYK model has “dual” description in which an extra spatial dimension, ζ , emerges. The curvature of this “emergent” spacetime is described by Einstein’s theory of general relativity



SYK and black holes



The BH entropy is proportional to the size of \mathbb{T}^2 , and hence the surface area of the black hole. Mapping to SYK applies when temperature $\ll 1/(\text{size of } \mathbb{T}^2)$.


SYK and black holes

Same long-time effective action for energy and number fluctuations, involving Schwarzian derivatives of time reparameterizations $f(\tau)$.



Einstein-Maxwell theory
+ cosmological constant

SYK and black holes



An extra spatial dimension emerges from quantum entanglement!


S. Sachdev, PRL **105**, 151602 (2010)

Tensor network of hierarchical entanglement

\vec{x}

D-dimensional space

space




depth of entanglement

String theory near
a "D-brane"

\vec{x}

D-dimensional
space



Emergent spatial direction
of SYK model or string theory

String theory near
a “D-brane”


\vec{x}

D-dimensional
space



Quantum entanglement
can lead to an emergent
spatial dimension

Emergent spatial direction
of SYK model or string theory




**Quantum
entanglement**

**Black
holes**

**Strange
metals**

**A "toy model" which is both a
strange metal and a black hole!**

Quantum Entanglement, Strange metals, and black holes



Subir Sachdev, Harvard University